

CRITICAL ZEROS OF THE RIEMANN ZETA-FUNCTION

H. M. BUI

ABSTRACT. In this unpublished note, we sketch an idea of using a three-piece mollifier to slightly improve the known percentages of zeros and simple zeros of the Riemann zeta-function on the critical line. This uses the recent result of Bettin *et al.* [2] on the twisted fourth moment of the Riemann zeta-function.

1. INTRODUCTION

Let $\zeta(s)$ be the Riemann zeta-function. Let $N(T)$ denote the number of zeros of $\zeta(s)$, $\rho = \beta + i\gamma$, with $0 < \gamma \leq T$ counted with multiplicity. Also let $N_0(T)$ denote the number of such critical zeros with $\beta = 1/2$, and $N_0^*(T)$ denote the number of such critical zeros with $\beta = 1/2$ and being simple. Define κ and κ^* by

$$\kappa = \liminf_{T \rightarrow \infty} \frac{N_0(T)}{N(T)}, \quad \kappa^* = \liminf_{T \rightarrow \infty} \frac{N_0^*(T)}{N(T)}.$$

Selberg [11] was the first to prove that a positive proportion of zeros lie on the critical line. Following the approach of Levinson [10] and the observation of Heath-Brown [9], it is now known that [3, 7]

$$\kappa > 0.410725 \quad \text{and} \quad \kappa^* > 0.40582. \quad (1)$$

Remark 1.1. Bui *et al.* [3] showed that $\kappa > 0.4105$ and $\kappa^* > 0.40582$. Shortly after that, Feng [8; version 1] put a paper on arXiv claiming that $\kappa > 0.4173$ and $\kappa^* > 0.4075$. Feng also used Levinson's method, but instead chose the mollifier as a sum of various pieces of different shapes $\sum_{n \leq T^{\vartheta_1}} + \dots + \sum_{n \leq T^{\vartheta_l}}$. At this point, Feng took $\vartheta_1 = \dots = \vartheta_l = 4/7 - \varepsilon$. The choice $\vartheta_1 = 4/7 - \varepsilon$ was from Conrey [4], but it was not clear why, say, $\vartheta_2 < 4/7$ is admissible. Conrey then emailed Feng requesting for the verification of that statement. Feng later agreed that that was a mistake, and subsequently replaced it with the second version. In this updated version [8; version 2], which is the same as the published one [7], he chose $\vartheta_1 = 4/7 - \varepsilon$ and $\vartheta_2 = \dots = \vartheta_l = 1/2 - \varepsilon$. With this Feng obtained $\kappa > 0.4128$, but no claim on the lower bound for κ^* . Feng still did not give any explanation as to why, say, the range $\vartheta_2 < 1/2$ is admissible. This is doubtful and can be problematic. Note that if one just applies the result of Balasubramanian *et al.* [1] on the twisted second moment of the Riemann zeta-function to, say, the cross term $\int |\zeta(1/2 + it)|^2 \sum_{n \leq T^{\vartheta_1}} \overline{\sum_{n \leq T^{\vartheta_2}}} dt$, then one needs $\vartheta_1 + \vartheta_2 < 1$. So without extra work, one can only take $\vartheta_1 = 4/7 - \varepsilon$ and $\vartheta_2 = \dots = \vartheta_l = 3/7 - \varepsilon$ in Feng's paper. This numerically leads to the above bound $\kappa > 0.410725$ in (1).

In this paper we shall prove

Theorem 1.1. *We have*

$$\kappa > 0.410918 \quad \text{and} \quad \kappa^* > 0.40589.$$

Remark 1.2. Rigourously speaking this is not yet a theorem. Here we assume a result on the twisted third moment of the Riemann zeta-function (see Theorem 5.1 for the precise

2010 *Mathematics Subject Classification.* 11M06, 11M26.

Key words and phrases. Riemann zeta-function, zeros on the critical line, mollifier method, moments.

statement). We leave that unproved. It is possible that the ideas of Bettin *et al.* [2] work in this context as well.

2. REDUCTION TO MEAN-VALUE THEOREMS

2.1. The mollifier. To get lower bounds for $N_0(T)$ and $N_0^*(T)$ it suffices to consider a certain mollified second moment of the Riemann zeta-function and its derivatives. This is well-known, so we shall simply state the conclusion.

Let $Q(x)$ be a real polynomial satisfying $Q(0) = 1$, and define

$$V(s) = Q\left(-\frac{1}{\mathcal{L}} \frac{d}{ds}\right) \zeta(s),$$

where for large T we denote

$$\mathcal{L} = \log T.$$

Suppose $\psi(s)$ is a “mollifier”. Littlewood’s lemma and the arithmetic-mean, geometric-mean inequality give

$$\kappa \geq 1 - \frac{1}{R} \log \left(\frac{1}{T} \int_1^T |V\psi(\sigma_0 + it)|^2 dt \right) + o(1), \quad (2)$$

where $\sigma_0 = 1/2 - R/\mathcal{L}$, and R is a bounded positive real number to be chosen later. Actually, by choosing $Q(x)$ to be a linear polynomial, we obtain a lower bound for the proportion of simple zeros, κ^* .

We choose a mollifier of the form

$$\psi(s) = \psi_1(s) + \psi_2(s) + \psi_3(s),$$

where ψ_1 , ψ_2 and ψ_3 are mollifiers of quite different shapes. Here ψ_1 is the usual mollifier

$$\psi_1(s) = \sum_{n \leq y_1} \frac{\mu(n) P_1\left(\frac{\log y_1/n}{\log y_1}\right) n^{\sigma_0-1/2}}{n^s}$$

and ψ_2 is the second piece mollifier used in [3]

$$\psi_2(s) = \chi(s + \tfrac{1}{2} - \sigma_0) \sum_{mn \leq y_2} \frac{\mu_2(m) P_2\left(\frac{\log y_2/mn}{\log y_2}\right) m^{\sigma_0-1/2} n^{1/2-\sigma_0}}{m^s n^{1-s}},$$

where $\mu_2(m)$ are the coefficients of $1/\zeta^2(s)$, and $P_1(x) = \sum_j a_j x^j$ and $P_2(x) = \sum_j b_j x^j$ are certain polynomials satisfying $P_1(0) = 0$ and $P_2(0) = P_2'(0) = P_2''(0) = 0$. For the third mollifier, we take¹

$$\psi_3(s) = \zeta(s + \tfrac{1}{2} - \sigma_0) \sum_{n \leq y_3} \frac{\mu_2(n) P_3\left(\frac{\log y_3/n}{\log y_3}\right) n^{\sigma_0-1/2}}{n^s},$$

where $P_3(x) = \sum_j c_j x^j$ is a certain polynomial satisfying $P_3(0) = P_3'(0) = P_3''(0) = P_3'''(0) = 0$. We also require the condition that $P_1(1) + P_3(1) = 1$ (see the below remark). Throughout the paper we denote $y_1 = T^{\vartheta_1}$, $y_2 = T^{\vartheta_2}$ and $y_3 = T^{\vartheta_3}$, where $0 < \vartheta_3 < \vartheta_2 < \vartheta_1 < 1$ (we shall see later what conditions are required on ϑ_1 , ϑ_2 and ϑ_3). Note that formally

$$\zeta(s) \sum_{n=1}^{\infty} \frac{\mu_2(n)}{n^s} = \frac{1}{\zeta(s)},$$

which explains why $\psi_3(s)$ may also be a useful choice of a mollifier.

¹It was originally an idea of K. Soundararajan to make use of the twisted fourth moment of the Riemann zeta-function and use a two-piece mollifier of the form $\psi_1 + \psi_3$. The author learned about this from Brian Conrey, Chris Hughes and K. Soundararajan. However, the implied proportion of critical zeros using this two-piece mollifier has never been worked out explicitly.

Remark 2.1. We can apply Levinson's method (see, for example, Appendix A of [5]) to our choice of $\psi = \psi_1 + \psi_2 + \psi_3$. It is important to note that when applying Littlewood's lemma, we need to estimate the integral on the right side of a rectangle. Assume that $\psi_1(s) + \psi_3(s)$ can be expressed as a Dirichlet series, $\psi_1(s) + \psi_3(s) = \sum_n a(n)n^{-s}$, that integral is negligible given that $a(1) = 1$. That is why we require $P_1(1) + P_3(1) = 1$.

2.2. A smoothing argument. It simplifies some calculations to smooth out the integral in (2). We introduce a smooth function $w(t)$ with the following properties

- i) $0 \leq w(t) \leq 1$ for all $t \in \mathbb{R}$,
- ii) w has compact support in $[T/4, 2T]$,
- iii) $w^{(j)}(t) \ll \Delta^{-j}$ for each $j = 0, 1, 2, \dots$, where $\Delta = T\mathcal{L}^{-1}$.

Theorem 2.1. Suppose $\vartheta_1 < 4/7$, $\vartheta_2 < 1/2$ and $\vartheta_3 < 1/4$. Then we have

$$\int_{-\infty}^{\infty} |V\psi(\sigma_0 + it)|^2 w(t) dt = c(P_1, P_2, P_3, Q, R, \vartheta_1, \vartheta_2, \vartheta_3) \widehat{w}(0) + O_{\varepsilon}(T\mathcal{L}^{-1+\varepsilon}), \quad (3)$$

where $c(P_1, P_2, P_3, Q, R, \vartheta_1, \vartheta_2, \vartheta_3) = c_1 + c_2 + c_3 + 2c_{12} + 2c_{23} + 2c_{31}$, and the c_i and c_{ij} are given below by (4)–(9).

2.3. Numerical evaluations. It is a standard exercise to deduce from Theorem 2.1 the unsmoothed version

$$\int_1^T |V\psi(\sigma_0 + it)|^2 dt = c(P_1, P_2, P_3, Q, R, \vartheta_1, \vartheta_2, \vartheta_3)T + O_{\varepsilon}(T\mathcal{L}^{-1+\varepsilon}).$$

Hence

$$\kappa \geq 1 - \frac{\log c(P_1, P_2, P_3, Q, R, \vartheta_1, \vartheta_2, \vartheta_3)}{R} + o(1).$$

Using Mathematica, with $R = 1.26$,

$$Q(x) = 0.49068 + 0.61077(1 - 2x) - 0.14199(1 - 2x)^3 + 0.04054(1 - 2x)^5,$$

$$P_1(x) = 0.83651x + 0.09758x^2 - 0.29393x^3 + 0.73372x^4 - 0.3753x^5,$$

$$P_2(x) = 0.0237x^3 - 0.00744x^4 + 0.00174x^5$$

and

$$P_3(x) = 0.00155x^4 - 0.00013x^5,$$

we get $\kappa > 0.410918$. To get $\kappa^* > 0.40589$ we take $R = 1.12$, $Q(x) = 1 - 1.03232x$,

$$P_1(x) = 0.82653x + 0.02626x^2 - 0.00774x^3 + 0.34803x^4 - 0.19371x^5,$$

$$P_2(x) = 0.0324x^3 - 0.00759x^4 + 0.00742x^5$$

and

$$P_3(x) = 0.00094x^4 - 0.00031x^5.$$

3. THE MEAN-VALUE RESULTS

Writing $\psi = \psi_1 + \psi_2 + \psi_3$ and opening the square, we get

$$\begin{aligned} \int |V\psi|^2 w &= \int |V\psi_1|^2 w + \int |V\psi_2|^2 w + \int |V\psi_3|^2 w \\ &\quad + 2\operatorname{Re}\left\{ \int |V|^2 \psi_1 \overline{\psi_2} w \right\} + 2\operatorname{Re}\left\{ \int |V|^2 \psi_2 \overline{\psi_3} w \right\} + 2\operatorname{Re}\left\{ \int |V|^2 \psi_3 \overline{\psi_1} w \right\} \\ &=: I_1 + I_2 + I_3 + 2\operatorname{Re}(I_{12}) + 2\operatorname{Re}(I_{23}) + 2\operatorname{Re}(I_{31}). \end{aligned}$$

We shall compute the integrals in turn. It turns out that I_{12} , I_{23} and I_{31} are asymptotically real.

3.1. The main terms. First we quote Theorem 2 of Conrey [4]. Conrey's theorem is for the unsmoothed version, but the following smoothed version follows easily from that.

Theorem 3.1 (Conrey). *Suppose $\vartheta_1 < 4/7$ and $P_1(0) = 0$. Then we have*

$$\int_{-\infty}^{\infty} |V\psi_1(\sigma_0 + it)|^2 w(t) dt = c_1 \widehat{w}(0) + O_{\varepsilon}(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$c_1 = P_1(1)^2 + \frac{1}{\vartheta_1} \int_0^1 \int_0^1 e^{2Rt} \left(Q(t)P_1'(u) + \vartheta_1 Q'(t)P_1(u) + \vartheta_1 RQ(t)P_1(u) \right)^2 dt du. \quad (4)$$

The terms I_{12} and I_2 are evaluated in [3].

Theorem 3.2 (Bui, Conrey and Young). *Suppose $\vartheta_2 < \vartheta_1 < 4/7$ and $P_1(0) = 0 = P_2(0) = P_2'(0)$. Then we have*

$$\int_{-\infty}^{\infty} |V|^2 \psi_1 \overline{\psi_2}(\sigma_0 + it) w(t) dt = c_{12} \widehat{w}(0) + O_{\varepsilon}(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$c_{12} = \frac{4\vartheta_2^2}{\vartheta_1^2} \frac{d^2}{dx_1 dx_2} \left[\int_0^1 \int_{\substack{t_1+t_2 \leq u \\ t_1, t_2 \geq 0}} e^{R(1-\vartheta_1(x_1-x_2)+\vartheta_2(t_1-t_2))} (1-u) Q(-\vartheta_1 x_1 + \vartheta_2 t_1) \right. \\ \left. Q(1 + \vartheta_1 x_2 - \vartheta_2 t_2) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_2}{\vartheta_1} (1-u) \right) P_2''(u - t_1 - t_2) dt_1 dt_2 du \right]_{x_1=x_2=0}. \quad (5)$$

Theorem 3.3 (Bui, Conrey and Young). *Suppose $\vartheta_2 < \frac{1}{2}$ and $P_2(0) = P_2'(0) = P_2''(0) = 0$. Then we have*

$$\int_{-\infty}^{\infty} |V\psi_2(\sigma_0 + it)|^2 w(t) dt = c_2 \widehat{w}(0) + O_{\varepsilon}(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$c_2 = \frac{2}{3\vartheta_2} \frac{d^4}{dx_1^2 dx_2^2} \left[\int_{[0,1]^4} e^{-\vartheta_2 R(x_1+x_2-t_1(x_1+u)-t_2(x_2+u))+2Rt_3(1+\vartheta_2(x_1+x_2-t_1(x_1+u)-t_2(x_2+u)))} \right. \\ \left(1 + \vartheta_2(x_1+x_2-t_1(x_1+u)-t_2(x_2+u)) \right) (x_1+u)(x_2+u)(1-u)^4 \\ Q \left(\vartheta_2(-x_1+t_2(x_2+u)) + t_3 \left(1 + \vartheta_2(x_1+x_2-t_1(x_1+u)-t_2(x_2+u)) \right) \right) \\ Q \left(\vartheta_2(-x_2+t_1(x_1+u)) + t_3 \left(1 + \vartheta_2(x_1+x_2-t_1(x_1+u)-t_2(x_2+u)) \right) \right) \\ \left. P_2''((x_1+u)(1-t_1)) P_2''((x_2+u)(1-t_2)) dt_1 dt_2 dt_3 du \right]_{x_1=x_2=0}. \quad (6)$$

We are left to evaluate I_3 , I_{23} and I_{31} , which are

$$I_3 = \int_{-\infty}^{\infty} |V(\sigma_0 + it) \zeta(\frac{1}{2} + it)|^2 \sum_{m \leq y_3} \frac{\mu_2(m) P_3(\frac{\log y_3/m}{\log y_3})}{m^{1/2+it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3(\frac{\log y_3/l}{\log y_3})}{l^{1/2-it}} w(t) dt, \\ I_{23} = \int_{-\infty}^{\infty} |V(\sigma_0 + it)|^2 \zeta(\frac{1}{2} + it) \sum_{mn \leq y_2} \frac{\mu_2(m) P_2(\frac{\log y_2/mn}{\log y_2})}{m^{1/2+it} n^{1/2-it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3(\frac{\log y_3/l}{\log y_3})}{l^{1/2-it}} w(t) dt$$

and

$$I_{31} = \int_{-\infty}^{\infty} |V(\sigma_0 + it)|^2 \zeta(\tfrac{1}{2} + it) \sum_{m \leq y_1} \frac{\mu(m) P_1(\frac{\log y_1/m}{\log y_1})}{m^{1/2-it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3(\frac{\log y_3/l}{\log y_3})}{l^{1/2+it}} w(t) dt.$$

Theorem 3.4. *Suppose $\vartheta_3 < 1/4$ and $P_3(0) = P'_3(0) = P''_3(0) = P'''_3(0) = 0$. Then we have*

$$I_3 = c_3 \widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$\begin{aligned} c_3 = & \frac{1}{12\vartheta_3^4} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[\int_{[0,1]^5} e^{-\vartheta_3 R(x_2+x_3) + Rt_1(1-t_4)(1+\vartheta_3(x_1+x_3)) + Rt_2(1-t_3)(1+\vartheta_3(x_2+x_4))} \right. \\ & e^{Rt_3(t_1+\vartheta_3(-x_1+x_2+(x_1+x_3)t_1)) + Rt_4(t_2+\vartheta_3(x_3-x_4+(x_2+x_4)t_2))} \\ & \left(t_1 - t_2 + \vartheta_3(-x_1+x_2+(x_1+x_3)t_1 - (x_2+x_4)t_2) \right) \\ & \left(t_1 - t_2 + \vartheta_3(-x_3+x_4+(x_1+x_3)t_1 - (x_2+x_4)t_2) \right) \\ & (1 + \vartheta_3(x_1+x_3))(1 + \vartheta_3(x_2+x_4))(1-u)^3 \\ & Q\left(-\vartheta_3 x_2 + t_2(1-t_3)(1 + \vartheta_3(x_2+x_4)) + t_3(t_1 + \vartheta_3(-x_1+x_2+(x_1+x_3)t_1))\right) \\ & Q\left(-\vartheta_3 x_3 + t_1(1-t_4)(1 + \vartheta_3(x_1+x_3)) + t_4(t_2 + \vartheta_3(x_3-x_4+(x_2+x_4)t_2))\right) \\ & \left. P_3(x_1+x_2+u)P_3(x_3+x_4+u)dt_1dt_2dt_3dt_4du \right]_{\underline{x}=0}. \end{aligned} \quad (7)$$

Theorem 3.5. *Suppose $\vartheta_2 < 1/2$, $\vartheta_3 < 1/4$ and $P_2(0) = P'_2(0) = P''_2(0) = 0 = P_3(0) = P'_3(0) = P''_3(0) = P'''_3(0)$. Then we have*

$$I_{23} = c_{23} \widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$\begin{aligned} c_{23} = & \frac{2}{3\vartheta_2^2} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_{[0,1]^5} e^{-R(\vartheta_2 x_1 + \vartheta_3 x_2) + Rt_1 t_2 (1-t_3-t_4 t_3)(\vartheta_2(1+x_1) - \vartheta_3(1-u))} \right. \\ & e^{Rt_3(1+t_4)(1+\vartheta_2 x_1 + \vartheta_3 x_2) + R(1-t_4)(\vartheta_3(x_2-x_3) - t_1(2t_2-1)(\vartheta_2(1+x_1) - \vartheta_3(1-u)))} \\ & \left(-\vartheta_3(x_2-x_3) + t_1(2t_2-1)(\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right. \\ & \left. + t_3(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2(\vartheta_2(1+x_1) - \vartheta_3(1-u))) \right) \\ & \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2(\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) t_1 \left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right)^2 (1-u)^3 \\ & Q\left(-\vartheta_3 x_2 + t_1 t_2(1-t_3 t_4)(\vartheta_2(1+x_1) - \vartheta_3(1-u)) + t_3 t_4(1 + \vartheta_2 x_1 + \vartheta_3 x_2) \right. \\ & \left. + (1-t_4)(\vartheta_3(x_2-x_3) - t_1(2t_2-1)(\vartheta_2(1+x_1) - \vartheta_3(1-u))) \right) \\ & Q\left(-\vartheta_2 x_1 + t_3(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2(\vartheta_2(1+x_1) - \vartheta_3(1-u))) \right) \\ & \left. P''_2\left((x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u))(1-t_1)\right) P_3(x_2+x_3+u)dt_1dt_2dt_3dt_4du \right]_{\underline{x}=0}. \end{aligned} \quad (8)$$

Theorem 3.6. Suppose $\vartheta_1 < 4/7$, $\vartheta_3 < 1/4$ and $P_1(0) = 0 = P_3(0) = P'_3(0)$. Then we have

$$I_{31} = c_{31}\widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

where

$$\begin{aligned} c_{31} = \frac{1}{\vartheta_1^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \Bigg[\int_{[0,1]^3} e^{-R(\vartheta_1 x_2 + \vartheta_3 x_3) + Rt_1(1+t_2)(1+\vartheta_1 x_1 + \vartheta_3 x_3) - \vartheta_1 Rt_2(x_1 - x_2)} \\ (-\vartheta_1(x_1 - x_2) + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3))(1 + \vartheta_1 x_1 + \vartheta_3 x_3)(1 - u) \\ Q(-\vartheta_1 x_2 + t_1 t_2(1 + \vartheta_1 x_1 + \vartheta_3 x_3) - \vartheta_1 t_2(x_1 - x_2)) Q(-\vartheta_3 x_3 + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3)) \\ P_1\left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1}(1 - u)\right) P_3(x_3 + u) dt_1 dt_2 du \Bigg]_{\underline{x}=0}. \end{aligned} \quad (9)$$

3.2. The shift parameters. Rather than working directly with $V(s)$, we shall instead consider the following three general integrals

$$\begin{aligned} I_3(\alpha, \beta, \gamma, \delta) = \int_{-\infty}^{\infty} \zeta\left(\frac{1}{2} + \alpha + it\right) \zeta\left(\frac{1}{2} + \beta + it\right) \zeta\left(\frac{1}{2} + \gamma - it\right) \zeta\left(\frac{1}{2} + \delta - it\right) \\ \sum_{m \leq y_3} \frac{\mu_2(m) P_3\left(\frac{\log y_3/m}{\log y_3}\right)}{m^{1/2+it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3\left(\frac{\log y_3/l}{\log y_3}\right)}{l^{1/2-it}} w(t) dt, \end{aligned} \quad (10)$$

$$\begin{aligned} I_{23}(\alpha, \beta, \gamma) = \int_{-\infty}^{\infty} \zeta\left(\frac{1}{2} + \alpha + it\right) \zeta\left(\frac{1}{2} + \beta + it\right) \zeta\left(\frac{1}{2} + \gamma - it\right) \\ \sum_{mn \leq y_2} \frac{\mu_2(m) P_2\left(\frac{\log y_2/mn}{\log y_2}\right)}{m^{1/2+it} n^{1/2-it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3\left(\frac{\log y_3/l}{\log y_3}\right)}{l^{1/2-it}} w(t) dt \end{aligned} \quad (11)$$

and

$$\begin{aligned} I_{31}(\alpha, \beta, \gamma) = \int_{-\infty}^{\infty} \zeta\left(\frac{1}{2} + \alpha + it\right) \zeta\left(\frac{1}{2} + \beta + it\right) \zeta\left(\frac{1}{2} + \gamma - it\right) \\ \sum_{m \leq y_1} \frac{\mu(m) P_1\left(\frac{\log y_1/m}{\log y_1}\right)}{m^{1/2-it}} \sum_{l \leq y_3} \frac{\mu_2(l) P_3\left(\frac{\log y_3/l}{\log y_3}\right)}{l^{1/2+it}} w(t) dt. \end{aligned} \quad (12)$$

Our main goal in the rest of the paper is to prove the following lemmas.

Lemma 3.1. Suppose $\vartheta_3 < 1/4$ and $P_3(0) = P'_3(0) = P''_3(0) = P'''_3(0) = 0$. Then we have

$$I_3(\alpha, \beta, \gamma, \delta) = c_3(\alpha, \beta, \gamma, \delta)\widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

uniformly for $\alpha, \beta, \gamma, \delta \ll \mathcal{L}^{-1}$, where $c_3(\alpha, \beta, \gamma, \delta)$ is given by

$$\begin{aligned} \frac{1}{12\vartheta_3^4} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \Bigg[\int_{[0,1]^5} y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (T y_3^{x_1+x_3})^{-(\alpha+\gamma)t_1} (T y_3^{x_2+x_4})^{-(\beta+\delta)t_2} \\ \left(T^{t_1-t_2} y_3^{-x_1+x_2+(x_1+x_3)t_1-(x_2+x_4)t_2} \right)^{-(\beta-\alpha)t_3} \left(T^{t_1-t_2} y_3^{-x_3+x_4+(x_1+x_3)t_1-(x_2+x_4)t_2} \right)^{-(\delta-\gamma)t_4} \\ \left(t_1 - t_2 + \vartheta_3(-x_1 + x_2 + (x_1 + x_3)t_1 - (x_2 + x_4)t_2) \right) \\ \left(t_1 - t_2 + \vartheta_3(-x_3 + x_4 + (x_1 + x_3)t_1 - (x_2 + x_4)t_2) \right) (1 + \vartheta_3(x_1 + x_3)) \\ (1 + \vartheta_3(x_2 + x_4))(1 - u)^3 P_3(x_1 + x_2 + u) P_3(x_3 + x_4 + u) dt_1 dt_2 dt_3 dt_4 du \Bigg]_{\underline{x}=0}. \end{aligned} \quad (13)$$

Remark 3.1. In the special case $\alpha = \beta = \gamma = \delta = 0$, Lemma 3.1 agrees with the mollified fourth moment of the Riemann zeta-function predicted by Conrey and Snaith using the ratios conjecture [6; Theorem 6.1].

Lemma 3.2. Suppose $\vartheta_2 < 1/2$, $\vartheta_3 < 1/4$ and $P_2(0) = P'_2(0) = P''_2(0) = 0 = P_3(0) = P'_3(0) = P''_3(0) = P'''_3(0)$. Then we have

$$I_{23}(\alpha, \beta, \gamma) = c_{23}(\alpha, \beta, \gamma)\widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

uniformly for $\alpha, \beta, \gamma \ll \mathcal{L}^{-1}$, where $c_{23}(\alpha, \beta, \gamma)$ is given by

$$\begin{aligned} & \frac{2}{3\vartheta_2^2} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_{[0,1]^5} y_2^{\gamma x_1 - (\alpha t_2 + \beta(1-t_2))t_1(1+x_1)} y_3^{\alpha x_2 + \beta x_3 + (\alpha t_2 + \beta(1-t_2))t_1(1-u)} \right. \\ & \quad (Ty_2^{x_1-t_1t_2(1+x_1)} y_3^{x_2+t_1t_2(1-u)})^{-(\alpha+\gamma)t_3} \\ & \quad \left(y_2^{t_1(2t_2-1)(1+x_1)} y_3^{-x_2+x_3-t_1(2t_2-1)(1-u)} (Ty_2^{x_1-t_1t_2(1+x_1)} y_3^{x_2+t_1t_2(1-u)})^{t_3} \right)^{-(\beta-\alpha)t_4} \\ & \quad \left(-\vartheta_3(x_2 - x_3) + t_1(2t_2 - 1)(\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right. \\ & \quad \left. + t_3 \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2 (\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) \right) \\ & \quad \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2 (\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) t_1 \left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right)^2 \\ & \quad \left. (1-u)^3 P''_2 \left((x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u))(1-t_1) \right) P_3(x_2 + x_3 + u) dt_1 dt_2 dt_3 du \right]_{\underline{x}=0}. \end{aligned} \quad (14)$$

Lemma 3.3. Suppose $\vartheta_1 < 4/7$, $\vartheta_3 < 1/4$ and $P_1(0) = 0 = P_3(0) = P'_3(0)$. Then we have

$$I_{31}(\alpha, \beta, \gamma) = c_{31}(\alpha, \beta, \gamma)\widehat{w}(0) + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),$$

uniformly for $\alpha, \beta, \gamma \ll \mathcal{L}^{-1}$, where $c_{31}(\alpha, \beta, \gamma)$ is given by

$$\begin{aligned} & \frac{1}{\vartheta_1^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_{[0,1]^3} y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (Ty_1^{x_1} y_3^{x_3})^{-(\alpha+\gamma)t_1} (y_1^{-x_1+x_2} (Ty_1^{x_1} y_3^{x_3})^{t_1})^{-(\beta-\alpha)t_2} \right. \\ & \quad \left(-\vartheta_1(x_1 - x_2) + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right) (1 + \vartheta_1 x_1 + \vartheta_3 x_3) \\ & \quad \left. (1-u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1}(1-u) \right) P_3(x_3 + u) dt_1 dt_2 du \right]_{\underline{x}=0}. \end{aligned} \quad (15)$$

We now prove that Theorems 3.4–3.6 follow from Lemmas 3.1–3.3, respectively. Let $I_\star$ denote either I_{23} or I_{31} . We first note that

$$I_3 = Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\beta}\right) Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\gamma}\right) I_3(\alpha, \beta, \gamma, \delta) \Big|_{\alpha=\delta=0, \beta=\gamma=-R/\mathcal{L}} \quad (16)$$

and

$$I_\star = Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\beta}\right) Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\gamma}\right) I_\star(\alpha, \beta, \gamma) \Big|_{\alpha=0, \beta=\gamma=-R/\mathcal{L}}. \quad (17)$$

We then argue that we can obtain either c_3 or $c_\star$ by applying the above differential operator to the corresponding $c_3(\alpha, \beta, \gamma, \delta)$ or $c_\star(\alpha, \beta, \gamma)$. Since $I_3(\alpha, \beta, \gamma, \delta)$, $I_\star(\alpha, \beta, \gamma)$, $c_3(\alpha, \beta, \gamma, \delta)$ and $c_\star(\alpha, \beta, \gamma)$ are holomorphic with respect to $\alpha, \beta, \gamma, \delta$ small, the derivatives appearing in (16) and (17) can be obtained as integrals of radii $\asymp \mathcal{L}^{-1}$ around the points $\alpha = \delta = 0$, $\beta = \gamma = -R/\mathcal{L}$, using Cauchy's integral formula. Since the error terms hold uniformly on

these contours, the same error terms that hold for $I_3(\alpha, \beta, \gamma, \delta)$ and $I_\star(\alpha, \beta, \gamma)$ also hold for I_3 and $I_\star$.

Next we check that applying the above differential operator to $c_3(\alpha, \beta, \gamma, \delta)$ and $c_\star(\alpha, \beta, \gamma)$ does indeed give c_3 and $c_\star$. Notice the formula

$$Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\beta}\right) X^{-\beta} = Q\left(\frac{\log X}{\mathcal{L}}\right) X^{-\beta}. \quad (18)$$

Using (18) and (15), we have

$$\begin{aligned} Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\beta}\right) Q\left(\frac{-1}{\mathcal{L}} \frac{d}{d\gamma}\right) c_{31}(0, \beta, \gamma) &= \frac{1}{\vartheta_1^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_{[0,1]^3} y_1^{\beta x_2} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-\gamma t_1} \right. \\ &\quad (y_1^{-x_1+x_2} (T y_1^{x_1} y_3^{x_3})^{t_1})^{-\beta t_2} \left(-\vartheta_1(x_1 - x_2) + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right) (1 + \vartheta_1 x_1 + \vartheta_3 x_3) \\ &\quad (1 - u) Q\left(-\vartheta_1 x_2 - \vartheta_1 t_2(x_1 - x_2) + t_1 t_2(1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right) \\ &\quad \left. Q\left(-\vartheta_3 x_3 + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right) P_1\left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1}(1 - u) \right) P_3(x_3 + u) dt_1 dt_2 du \right]_{\underline{x}=0}. \end{aligned}$$

Setting $\beta = \gamma = -R/\mathcal{L}$ and simplifying gives (9). A similar argument produces (8) from (14) and produces (7) from (13).

We prove Lemma 3.3 in Section 5, Lemma 3.2 in Section 6 and Lemma 3.1 in Section 7.

4. VARIOUS LEMMAS

4.1. The Euler-Maclaurin formula. The following two lemmas are easy consequences of the Euler-Maclaurin formula, see [3; Lemmas 4.4 and 4.6].

Lemma 4.1. *Suppose $y_2 \leq y_1$, $|z| \ll (\log y_2)^{-1}$, and that f_1 and f_2 are smooth functions. Then we have*

$$\begin{aligned} \sum_{n \leq y_2} \frac{d_k(n)}{n} \left(\frac{y_2}{n}\right)^z f_1\left(\frac{\log y_1/n}{\log y_1}\right) f_2\left(\frac{\log y_2/n}{\log y_2}\right) \\ = \frac{(\log y_2)^k}{(k-1)!} \int_0^1 y_2^{zu} (1-u)^{k-1} f_1\left(1 - \frac{(1-u)\log y_2}{\log y_1}\right) f_2(u) du + O((\log y_2)^{k-1}). \end{aligned}$$

Lemma 4.2. *Suppose $-1 \leq \sigma \leq 0$. Then we have*

$$\sum_{n \leq y_1} \frac{d_k(n)}{n} \left(\frac{y_1}{n}\right)^\sigma \ll (\log y_1)^{k-1} \min\{|\sigma|^{-1}, \log y_1\}.$$

4.2. Mellin pairs. By convention, we set $P_j(x) = 0$ for $j = 1, 2, 3$ and $x \leq 0$. Note that with this definition we have

$$P_1\left(\frac{\log y_1/n}{\log y_1}\right) = \sum_j \frac{a_j j!}{(\log y_1)^j} \frac{1}{2\pi i} \int_{(1)} \left(\frac{y_1}{n}\right)^u \frac{du}{u^{j+1}} \quad (19)$$

for all n . Similar expressions holds for $P_2\left(\frac{\log y_2/n}{\log y_2}\right)$ and $P_3\left(\frac{\log y_3/n}{\log y_3}\right)$.

5. PROOF OF LEMMA 3.3

5.1. Reduction to a contour integral. We shall use the following unproved twisted third moment of the Riemann zeta-function.

Theorem 5.1. *Suppose H and K satisfy $H \leq T^{4/7-\varepsilon}$ and $K \leq T^{1/4-\varepsilon}$, or $HK \leq T^{3/4-\varepsilon}$ and $K \leq T^{1/2-\varepsilon^2}$. Then we have*

$$\begin{aligned} \sum_{\substack{h \leq H \\ k \leq K}} \frac{a_h \bar{a}_k}{\sqrt{hk}} \int_{-\infty}^{\infty} \zeta\left(\frac{1}{2} + \alpha + it\right) \zeta\left(\frac{1}{2} + \beta + it\right) \zeta\left(\frac{1}{2} + \gamma - it\right) \left(\frac{h}{k}\right)^{it} w(t) dt \\ = \sum_{\substack{h \leq H \\ k \leq K}} \frac{a_h \bar{a}_k}{\sqrt{hk}} \int_{-\infty}^{\infty} w(t) \left\{ Z_{\alpha, \beta, \gamma}(h, k) + \left(\frac{t}{2\pi}\right)^{-(\alpha+\gamma)} Z_{-\gamma, \beta, -\alpha}(h, k) \right. \\ \left. + \left(\frac{t}{2\pi}\right)^{-(\beta+\gamma)} Z_{\alpha, -\gamma, -\beta}(h, k) \right\} dt + O_{\varepsilon}(T^{1-\varepsilon}) \end{aligned}$$

uniformly for $\alpha, \beta, \gamma \ll \mathcal{L}^{-1}$, where

$$Z_{\alpha, \beta, \gamma}(h, k) = \sum_{kab=hc} \frac{1}{a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}}.$$

Recall that $I_{31}(\alpha, \beta, \gamma)$ is defined by (12). We have

$$I_{31}(\alpha, \beta, \gamma) = I_{31}^1 + I_{31}^2 + I_{31}^3 + O_{\varepsilon}(T^{1-\varepsilon}),$$

where

$$I_{31}^1 = \widehat{w}(0) \sum_{l, m} \frac{\mu(m) \mu_2(l) P_1\left(\frac{\log y_1/m}{\log y_1}\right) P_3\left(\frac{\log y_3/l}{\log y_3}\right)}{\sqrt{lm}} \sum_{lab=mc} \frac{1}{a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}},$$

I_{31}^2 is obtained by multiplying I_{31}^1 with $T^{-(\alpha+\gamma)}$ and changing the shifts $\alpha \leftrightarrow -\gamma$, $\gamma \leftrightarrow -\alpha$, and I_{31}^3 is obtained by multiplying I_{31}^1 with $T^{-(\beta+\gamma)}$ and changing the shifts $\beta \leftrightarrow -\gamma$, $\gamma \leftrightarrow -\beta$.

We shall first work on I_{31}^1 . In view of (19) we get

$$\begin{aligned} I_{31}^1 &= \widehat{w}(0) \sum_{i, j} \frac{a_i c_j i! j!}{(\log y_1)^i (\log y_3)^j} \left(\frac{1}{2\pi i}\right)^2 \int_{(1)} \int_{(1)} y_1^u y_3^v \\ &\quad \sum_{lab=mc} \frac{\mu(m) \mu_2(l)}{m^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}. \end{aligned}$$

The Euler product implies that

$$\begin{aligned} \sum_{lab=mc} \frac{\mu(m) \mu_2(l)}{m^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}} \\ = A(\alpha, \beta, \gamma, u, v) \frac{\zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \zeta(1+u+v)^2}{\zeta(1+\alpha+u) \zeta(1+\beta+u) \zeta(1+\gamma+v)^2}, \end{aligned} \quad (20)$$

where $A(\alpha, \beta, \gamma, u, v)$ is an arithmetical factor converging absolutely in a product of half-planes containing the origin. Hence

$$I_{31}^1 = \widehat{w}(0) \zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \sum_{i, j} \frac{a_i c_j i! j!}{(\log y_1)^i (\log y_3)^j} J_{i, j}, \quad (21)$$

where

$$J_{i, j} = \left(\frac{1}{2\pi i}\right)^2 \int_{(1)} \int_{(1)} y_1^u y_3^v \frac{A(\alpha, \beta, \gamma, u, v) \zeta(1+u+v)^2}{\zeta(1+\alpha+u) \zeta(1+\beta+u) \zeta(1+\gamma+v)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}.$$

²Theorem 5.1 is likely to hold for larger ranges of H and K . These conditions, however, suffice for our purposes.

Using the Dirichlet series for $\zeta(1+u+v)^2$ and reversing the order of summation and integration, we obtain

$$J_{i,j} = \sum_{n \leq y_3} \frac{d(n)}{n} \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} \left(\frac{y_1}{n} \right)^u \left(\frac{y_3}{n} \right)^v \frac{A(\alpha, \beta, \gamma, u, v)}{\zeta(1+\alpha+u)\zeta(1+\beta+u)\zeta(1+\gamma+v)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}. \quad (22)$$

Note that here we are able to restrict the sum over n to $n \leq y_3$ by moving the v -integral far to the right. We now move the contours of integration to $\text{Re}(u) = \text{Re}(v) \asymp \mathcal{L}^{-1}$. Bounding the integrals trivially shows that $J_{i,j} \ll \mathcal{L}^{i+j-2}$. Hence from the Taylor series $A(\alpha, \beta, \gamma, u, v) = A(0, 0, 0, 0, 0) + O(\mathcal{L}^{-1}) + O(|u| + |v|)$, we can replace $A(\alpha, \beta, \gamma, u, v)$ by $A(0, 0, 0, 0, 0)$ in $J_{i,j}$ with an error of size $O(\mathcal{L}^{i+j-3})$. By letting $\alpha = \beta = \gamma = u = v = s$ in (20), it is easy to verify that $A(0, 0, 0, 0, 0) = 1$. The u and v variables in (22) are now separated so that

$$J_{i,j} = \sum_{n \leq y_3} \frac{d(n)}{n} M_i(\alpha, \beta) M_j(\gamma) + O(\mathcal{L}^{i+j-3}), \quad (23)$$

where

$$M_i(\alpha, \beta) = \frac{1}{2\pi i} \int_{(\mathcal{L}^{-1})} \left(\frac{y_1}{n} \right)^u \frac{1}{\zeta(1+\alpha+u)\zeta(1+\beta+u)} \frac{du}{u^{i+1}}$$

and

$$M_j(\gamma) = \frac{1}{2\pi i} \int_{(\mathcal{L}^{-1})} \left(\frac{y_3}{n} \right)^v \frac{1}{\zeta(1+\gamma+v)^2} \frac{dv}{v^{j+1}}.$$

The expression $M_i(\alpha, \beta)$ was evaluated in [3; Lemma 5.7]

$$M_i(\alpha, \beta) = \frac{1}{i! (\log y_1)^2} \frac{d^2}{dx_1 dx_2} \left[y_1^{\alpha x_1 + \beta x_2} \left((x_1 + x_2) \log y_1 + \log y_1/n \right)^i \right]_{x_1=x_2=0} + O(\mathcal{L}^{i-3}). \quad (24)$$

We evaluate $M_j(\gamma)$ with the following lemma.

Lemma 5.1. *Suppose $j \geq 2$. Then for some $\nu \asymp (\log \log y_3)^{-1}$ we have*

$$M_j(\gamma) = \frac{1}{j! (\log y_3)^2} \frac{d^2}{dx_3^2} \left[y_3^{\gamma x_3} (x_3 \log y_3 + \log y_3/n)^j \right]_{x_3=0} + O(\mathcal{L}^{j-3}) + O_\varepsilon \left(\left(\frac{y_3}{n} \right)^{-\nu} \mathcal{L}^\varepsilon \right).$$

Proof. Let $Y = o(T)$ be a large parameter to be chosen later. By Cauchy's theorem, $M_j(\gamma)$ is equal to the residue at $v = 0$ plus integrals over the line segments $\mathcal{C}_1 = \{v = \mathcal{L}^{-1} + it, t \in \mathbb{R}, |t| \geq Y\}$, $\mathcal{C}_2 = \{v = \sigma \pm iY, -\frac{c}{\log Y} \leq \sigma \leq \mathcal{L}^{-1}\}$ and $\mathcal{C}_3 = \{v = -\frac{c}{\log Y} + it, |t| \leq Y\}$, where c is some fixed positive constant such that $\zeta(1+\gamma+v)$ has no zeros in the region on the right hand side of the contour determined by the $\mathcal{C}_j$. Furthermore, we require that for such c we have $1/\zeta(\sigma + it) \ll \log(2 + |t|)$ in this region (see [12; Theorem 3.11]). Then the integral over $\mathcal{C}_1$ is

$$\ll (\log Y)^2 / Y^j \ll_\varepsilon Y^{-2+\varepsilon},$$

since $j \geq 2$. The integral over $\mathcal{C}_2$ is

$$\ll (\log Y) / Y^{j+1} \ll_\varepsilon Y^{-3+\varepsilon}.$$

Finally, the contribution from $\mathcal{C}_3$ is

$$\ll (\log Y)^j \left(\frac{y_3}{n} \right)^{-c/\log Y} \ll_\varepsilon \left(\frac{y_3}{n} \right)^{-c/\log Y} Y^\varepsilon.$$

Choosing $Y \asymp \mathcal{L}$ gives an error so far of size $O_\varepsilon((y_3/n)^{-\nu} \mathcal{L}^\varepsilon) + O_\varepsilon(\mathcal{L}^{-2+\varepsilon})$.

For the residue at $v = 0$, we write this as

$$\frac{1}{2\pi i} \oint \left(\frac{y_3}{n} \right)^v \frac{1}{\zeta(1+\gamma+v)^2} \frac{dv}{v^{j+1}},$$

where the contour is a circle of radius $\asymp \mathcal{L}^{-1}$ around the origin. This integral is trivially bounded by $O(\mathcal{L}^{j-2})$, hence by taking the first term in the Taylor series of $\zeta(1+\gamma+v)$ we get

$$\text{Res}_{v=0} = \frac{\gamma^2(\log y_3/n)^j}{j!} + \frac{2\gamma(\log y_3/n)^{j-1}}{(j-1)!} + \frac{(\log y_3/n)^{j-2}}{(j-2)!} + O(\mathcal{L}^{j-3}).$$

The above main term can be written in a compact form

$$\frac{1}{j!(\log y_3)^2} \frac{d^2}{dx_3^2} \left[y_3^{\gamma x_3} (x_3 \log y_3 + \log y_3/n)^j \right]_{x_3=0}$$

and the lemma follows. $\square$

In view of (23), (24) and Lemma 5.1 we get

$$\begin{aligned} J_{i,j} &= \frac{(\log y_1)^{i-2}(\log y_3)^{j-2}}{i!j!} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} \right. \\ &\quad \left. \sum_{n \leq y_3} \frac{d(n)}{n} \left(x_1 + x_2 + \frac{\log y_1/n}{\log y_1} \right)^i \left(x_3 + \frac{\log y_3/n}{\log y_3} \right)^j \right]_{\underline{x}=0} \\ &\quad + O_\varepsilon \left(\mathcal{L}^{i-2+\varepsilon} \sum_{n \leq y_3} \frac{d(n)}{n} \left(\frac{y_3}{n} \right)^{-\nu} \right) + O(\mathcal{L}^{i+j-3}). \end{aligned}$$

Using Lemmas 4.1 and 4.2 this is equal to

$$\begin{aligned} &\frac{(\log y_1)^{i-2}(\log y_3)^j}{i!j!} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} \right. \\ &\quad \left. \int_0^1 (1-u) \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1-u) \right)^i (x_3 + u)^j du \right]_{\underline{x}=0} + O_\varepsilon(\mathcal{L}^{i-1+\varepsilon}) + O(\mathcal{L}^{i+j-3}). \end{aligned}$$

As $j \geq 2$, putting this back to (21) we get

$$\begin{aligned} I_{31}^1 &= \frac{\widehat{w}(0)}{(\log y_1)^2} \zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \frac{d^4}{dx_1 dx_2 dx_3^2} \left[y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} \right. \\ &\quad \left. \int_0^1 (1-u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1-u) \right) P_3(x_3 + u) du \right]_{\underline{x}=0} + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}). \end{aligned}$$

5.2. Deduction of Lemma 3.3. Combining I_{31}^1 , I_{31}^2 and I_{31}^3 we have

$$\begin{aligned} I_{31} &= \frac{\widehat{w}(0)}{(\log y_1)^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_0^1 U_1(\underline{x}) (1-u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1-u) \right) P_3(x_3 + u) du \right]_{\underline{x}=0} \\ &\quad + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}), \end{aligned}$$

where

$$U_1 = \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3}}{(\alpha + \gamma)(\beta + \gamma)} - \frac{T^{-(\alpha + \gamma)} y_1^{-\gamma x_1 + \beta x_2} y_3^{-\alpha x_3}}{(\alpha + \gamma)(\beta - \alpha)} - \frac{T^{-(\beta + \gamma)} y_1^{\alpha x_1 - \gamma x_2} y_3^{-\beta x_3}}{(\alpha - \beta)(\beta + \gamma)}.$$

Using the identity

$$\frac{1}{(\alpha + \gamma)(\beta + \gamma)} = \frac{1}{(\alpha + \gamma)(\beta - \alpha)} + \frac{1}{(\alpha - \beta)(\beta + \gamma)}$$

and the integral formula

$$\frac{1 - z^{-(\alpha + \beta)}}{\alpha + \beta} = (\log z) \int_0^1 z^{-(\alpha + \beta)t} dt, \quad (25)$$

we can write

$$\begin{aligned}
U_1 &= \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} - T^{-(\alpha + \gamma)} y_1^{-\gamma x_1 + \beta x_2} y_3^{-\alpha x_3}}{(\alpha + \gamma)(\beta - \alpha)} - \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} - T^{-(\beta + \gamma)} y_1^{\alpha x_1 - \gamma x_2} y_3^{-\beta x_3}}{(\beta - \alpha)(\beta + \gamma)} \\
&= \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3}}{\beta - \alpha} \left(\frac{1 - (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)}}{\alpha + \gamma} \right) - \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3}}{\beta - \alpha} \left(\frac{1 - (T y_1^{x_2} y_3^{x_3})^{-(\beta + \gamma)}}{\beta + \gamma} \right) \\
&= \frac{\mathcal{L}}{\beta - \alpha} (1 + \vartheta_1 x_1 + \vartheta_3 x_3) y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} \int_0^1 (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)t_1} dt_1 \\
&\quad - \frac{\mathcal{L}}{\beta - \alpha} (1 + \vartheta_1 x_2 + \vartheta_3 x_3) y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} \int_0^1 (T y_1^{x_2} y_3^{x_3})^{-(\beta + \gamma)t_1} dt_1.
\end{aligned}$$

Hence

$$\begin{aligned}
I_{31} &= \frac{\widehat{w}(0)\mathcal{L}}{(\beta - \alpha)(\log y_1)^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_0^1 \int_0^1 y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)t_1} (1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right. \\
&\quad \left. (1 - u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1 - u) \right) P_3(x_3 + u) dt_1 du \right]_{\underline{x}=0} \\
&\quad - \frac{\widehat{w}(0)\mathcal{L}}{(\beta - \alpha)(\log y_1)^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_0^1 \int_0^1 y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (T y_1^{x_2} y_3^{x_3})^{-(\beta + \gamma)t_1} (1 + \vartheta_1 x_2 + \vartheta_3 x_3) \right. \\
&\quad \left. (1 - u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1 - u) \right) P_3(x_3 + u) dt_1 du \right]_{\underline{x}=0} + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}).
\end{aligned}$$

Changing the roles of the variables x_1 and x_2 in the second term we obtain

$$\begin{aligned}
I_{31} &= \frac{\widehat{w}(0)\mathcal{L}}{(\log y_1)^2} \frac{d^4}{dx_1 dx_2 dx_3^2} \left[\int_0^1 \int_0^1 V_1(\underline{x}) (1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right. \\
&\quad \left. (1 - u) P_1 \left(x_1 + x_2 + 1 - \frac{\vartheta_3}{\vartheta_1} (1 - u) \right) P_3(x_3 + u) dt_1 du \right]_{\underline{x}=0} + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}),
\end{aligned} \tag{26}$$

where

$$V_1 = \frac{y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)t_1} - y_1^{\alpha x_2 + \beta x_1} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-(\beta + \gamma)t_1}}{\beta - \alpha}.$$

Using (25) again we have

$$\begin{aligned}
V_1 &= y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)t_1} \left(\frac{1 - (y_1^{-x_1 + x_2} (T y_1^{x_1} y_3^{x_3})^{t_1})^{-(\beta - \alpha)}}{\beta - \alpha} \right) \\
&= \mathcal{L} \left(-\vartheta_1(x_1 - x_2) + t_1(1 + \vartheta_1 x_1 + \vartheta_3 x_3) \right) y_1^{\alpha x_1 + \beta x_2} y_3^{\gamma x_3} (T y_1^{x_1} y_3^{x_3})^{-(\alpha + \gamma)t_1} \\
&\quad \int_0^1 (y_1^{-x_1 + x_2} (T y_1^{x_1} y_3^{x_3})^{t_1})^{-(\beta - \alpha)t_2} dt_2.
\end{aligned}$$

Lemma 3.3 follows from this and (26).

6. PROOF OF LEMMA 3.2

6.1. Reduction to a contour integral. Recall that $I_{23}(\alpha, \beta, \gamma)$ is defined by (11). We have

$$I_{23}(\alpha, \beta, \gamma) = I_{23}^1 + I_{23}^2 + I_{23}^3,$$

where

$$I_{23}^1 = \widehat{w}(0) \sum_{l, m, n} \frac{\mu_2(m) \mu_2(l) P_2\left(\frac{\log y_2 / mn}{\log y_2}\right) P_3\left(\frac{\log y_3 / l}{\log y_3}\right)}{\sqrt{lmn}} \sum_{mab=ln c} \frac{1}{a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}},$$

I_{23}^2 is obtained by multiplying I_{23}^1 with $T^{-(\alpha+\gamma)}$ and changing the shifts $\alpha \leftrightarrow -\gamma$, $\gamma \leftrightarrow -\alpha$, and I_{23}^3 is obtained by multiplying I_{23}^1 with $T^{-(\beta+\gamma)}$ and changing the shifts $\beta \leftrightarrow -\gamma$, $\gamma \leftrightarrow -\beta$.

We first work on I_{23}^1 . In view of (19) we get

$$I_{23}^1 = \widehat{w}(0) \sum_{i,j} \frac{b_i c_j i! j!}{(\log y_2)^i (\log y_3)^j} \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} y_2^u y_3^v \sum_{mab=lnc} \frac{\mu_2(m) \mu_2(l)}{(mn)^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}.$$

The arithmetical sum is

$$\begin{aligned} & \sum_{mab=lnc} \frac{\mu_2(m) \mu_2(l)}{(mn)^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma}} \\ &= B(\alpha, \beta, \gamma, u, v) \frac{\zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \zeta(1+\alpha+u) \zeta(1+\beta+u) \zeta(1+u+v)^4}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+2u)^2}, \end{aligned} \quad (27)$$

where $B(\alpha, \beta, \gamma, u, v)$ is an arithmetical factor converging absolutely in a product of half-planes containing the origin. So

$$I_{23}^1 = \widehat{w}(0) \zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \sum_{i,j} \frac{b_i c_j i! j!}{(\log y_2)^i (\log y_3)^j} K_{i,j}, \quad (28)$$

where

$$K_{i,j} = \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} y_2^u y_3^v \frac{B(\alpha, \beta, \gamma, u, v) \zeta(1+\alpha+u) \zeta(1+\beta+u) \zeta(1+u+v)^4}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+2u)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}.$$

Using the Dirichlet series for $\zeta(1+u+v)^4$ and changing the order of summation and integration, we have

$$\begin{aligned} K_{i,j} &= \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} \left(\frac{y_2}{n} \right)^u \left(\frac{y_3}{n} \right)^v \\ & \quad \frac{B(\alpha, \beta, \gamma, u, v) \zeta(1+\alpha+u) \zeta(1+\beta+u)}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+2u)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}. \end{aligned} \quad (29)$$

Note that here we are able to restrict the sum over n to $n \leq y_3$ by moving the v -integral far to the right. We now move the contours of integration to $\text{Re}(u) = \text{Re}(v) \asymp \mathcal{L}^{-1}$. Bounding the integrals trivially shows that $K_{i,j} \ll \mathcal{L}^{i+j-2}$. Hence from the Taylor series $B(\alpha, \beta, \gamma, u, v) = B(0, 0, 0, 0, 0) + O(\mathcal{L}^{-1}) + O(|u| + |v|)$, we can replace $B(\alpha, \beta, \gamma, u, v)$ by $B(0, 0, 0, 0, 0)$ in $K_{i,j}$ with an error of size $O(\mathcal{L}^{i+j-3})$. By letting $\alpha = \beta = \gamma = u = v = s$ in (27), it is easy to verify that $B(0, 0, 0, 0, 0) = 1$. The u and v variables in (29) are now separated so that

$$K_{i,j} = \sum_{n \leq y_3} \frac{d_4(n)}{n} N_i(\alpha, \beta, \gamma) N_j(\alpha, \beta) + O(\mathcal{L}^{i+j-3}), \quad (30)$$

where

$$N_i(\alpha, \beta, \gamma) = \frac{1}{2\pi i} \int_{(\mathcal{L}^{-1})} \left(\frac{y_2}{n} \right)^u \frac{\zeta(1+\alpha+u) \zeta(1+\beta+u)}{\zeta(1+\gamma+u)^2 \zeta(1+2u)^2} \frac{du}{u^{i+1}}$$

and

$$N_j(\alpha, \beta) = \frac{1}{2\pi i} \int_{(\mathcal{L}^{-1})} \left(\frac{y_3}{n} \right)^v \frac{1}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2} \frac{dv}{v^{j+1}}. \quad (31)$$

We evaluate $N_i(\alpha, \beta, \gamma)$ and $N_j(\alpha, \beta)$ with the following lemmas.

Lemma 6.1. *Suppose $i \geq 3$. Then we have*

$$N_i(\alpha, \beta, \gamma) = \frac{4}{(i-2)!(\log y_2)^2} \frac{d^2}{dx_1^2} \left[\int_0^1 \int_0^1 y_2^{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 x_1} \left(\frac{y_2}{n} \right)^{-(\alpha(1-t_2) + \beta t_2) t_1} (1-t_1)^{i-2} t_1 \left(x_1 \log y_2 + (\log y_2/n) \right)^i dt_1 dt_2 \right]_{x_1=0} + O(\mathcal{L}^{i-3}).$$

Proof. An argument on the level of the prime number theorem shows that $N_i(\alpha, \beta, \gamma)$ is equal to the residue at $u = 0$ plus an error of size $(\log y_2/n)^{-B}$ for any $B > 0$. Since $n \leq y_3$, we have $(\log y_2/n) \geq \log(y_2/y_3) = (\vartheta_2 - \vartheta_3)\mathcal{L}$ so that this error is negligible.

We write the residue at $u = 0$ as

$$\frac{1}{2\pi i} \oint q^u \frac{\zeta(1+\alpha+u)\zeta(1+\beta+u)}{\zeta(1+\gamma+u)^2\zeta(1+2u)^2} \frac{du}{u^{i+1}},$$

where $q = y_2/n$ and the contour is a circle of radius $\asymp \mathcal{L}^{-1}$ around the origin. This integral is trivially bounded by $O(\mathcal{L}^{i-2})$. Hence by taking the first terms in the Taylor series of the zeta-functions we have

$$N_i(\alpha, \beta, \gamma) = \frac{4}{2\pi i} \oint q^u \frac{(\gamma+u)^2}{(\alpha+u)(\beta+u)} \frac{du}{u^{i-1}} + O(\mathcal{L}^{i-3}).$$

We next use the identity

$$(\gamma+u)^2 = \frac{1}{(\log y_2)^2} \frac{d^2}{dx_1^2} y_2^{(\gamma+u)x_1} \Big|_{x_1=0},$$

to write

$$N_i(\alpha, \beta, \gamma) = \frac{4}{(\log y_2)^2} \frac{d^2}{dx_1^2} y_2^{\gamma x_1} N(x_1) \Big|_{x_1=0},$$

where

$$N(x_1) = \frac{1}{2\pi i} \oint (y_2^{x_1} q)^u \frac{1}{(\alpha+u)(\beta+u)} \frac{du}{u^{i-1}}.$$

Taking the power series gives

$$N(x_1) = \sum_{k \geq 0} \frac{(x_1 \log y_2 + \log q)^k}{k!} \frac{1}{2\pi i} \oint \frac{u^{k-i+1}}{(\alpha+u)(\beta+u)} du.$$

As there are three poles inside the contour, it is slightly easier to compute the residue at infinity. In other words, changing the variable $u \rightarrow 1/u$ yields

$$N(x_1) = \sum_{k \geq 0} \frac{(x_1 \log y_2 + \log q)^k}{k!} \frac{1}{2\pi i} \oint \frac{u^{i-k-1}}{(1+\alpha u)(1+\beta u)} du.$$

In view of the power series of $(1+\alpha u)^{-1}$ and $(1+\beta u)^{-1}$, we have

$$N(x_1) = \sum_{k \geq 0} \frac{(x_1 \log y_2 + \log q)^k}{k!} \sum_{l_1, l_2 \geq 0} (-\alpha)^{l_1} (-\beta)^{l_2} \frac{1}{2\pi i} \oint u^{i+l_1+l_2-k-1} du.$$

The integral picks out the terms $k = i + l_1 + l_2$, giving

$$N(x_1) = (x_1 \log y_2 + \log q)^i \sum_{l_1, l_2 \geq 0} \frac{(-\alpha)^{l_1} (-\beta)^{l_2} (x_1 \log y_2 + \log q)^{l_1+l_2}}{(i+l_1+l_2)!}. \quad (32)$$

We now separate the variables l_1, l_2 and i by using the standard beta function and its integral representation

$$\frac{1}{(i+l_1+l_2)!} = B(i-1, l_1+l_2+2) \frac{1}{(i-2)!(l_1+l_2+1)!}$$

$$\begin{aligned}
&= B(i-1, l_1+l_2+2)B(l_1+1, l_2+1)\frac{1}{(i-2)!l_1!l_2!} \\
&= \frac{1}{(i-2)!l_1!l_2!} \int_0^1 \int_0^1 (1-t_1)^{i-2} t_1^{l_1+l_2+1} (1-t_2)^{l_1} t_2^{l_2} dt_1 dt_2.
\end{aligned}$$

Putting this into (32) we obtain

$$N(x_1) = \frac{(x_1 \log y_2 + \log q)^i}{(i-2)!} \int_0^1 \int_0^1 (1-t_1)^{i-2} t_1 e^{-\alpha(x_1 \log y_2 + \log q)t_1(1-t_2) - \beta(x_1 \log y_2 + \log q)t_1 t_2} dt_1 dt_2,$$

and the lemma follows. $\square$

Lemma 6.2. *Suppose $j \geq 4$. Then for some $\nu \asymp (\log \log y_3)^{-1}$ we have*

$$\begin{aligned}
N_j(\alpha, \beta) &= \frac{1}{j!(\log y_3)^4} \frac{d^4}{dx_2^2 dx_3^2} \left[y_3^{\alpha x_2 + \beta x_3} \left((x_2 + x_3) \log y_3 + \log y_3/n \right)^j \right]_{x_2=x_3=0} \\
&\quad + O(\mathcal{L}^{j-5}) + O\left(\left(\frac{y_3}{n} \right)^{-\nu} \mathcal{L}^\varepsilon \right).
\end{aligned}$$

Proof. Similarly to Lemma 5.1, $N_j(\alpha, \beta)$ equals the residue at $v = 0$ plus an error of size $O_\varepsilon((y_3/n)^{-\nu} \mathcal{L}^\varepsilon) + O_\varepsilon(\mathcal{L}^{-2+\varepsilon})$.

For the residue at $v = 0$, we write this as

$$\frac{1}{2\pi i} \oint \left(\frac{y_3}{n} \right)^v \frac{1}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2} \frac{dv}{v^{j+1}},$$

where the contour is a circle of radius $\asymp \mathcal{L}^{-1}$ around the origin. This integral is trivially bounded by $O(\mathcal{L}^{j-4})$. Hence by taking the first terms in the Taylor series of the zeta-functions we have

$$\text{Res}_{v=0} = \frac{1}{2\pi i} \oint \left(\frac{y_3}{n} \right)^v (\alpha+v)^2 (\beta+v)^2 \frac{dv}{v^{j+1}} + O(\mathcal{L}^{j-5}).$$

The above integral can be written in a compact form as

$$\frac{1}{j!(\log y_3)^4} \frac{d^4}{dx_2^2 dx_3^2} \left[y_3^{\alpha x_2 + \beta x_3} \left((x_2 + x_3) \log y_3 + \log y_3/n \right)^j \right]_{x_2=x_3=0},$$

and the lemma follows. $\square$

In view of (30) and Lemmas 6.1 and 6.2 we get

$$\begin{aligned}
K_{i,j} &= \frac{4(\log y_2)^{i-2} (\log y_3)^{j-4}}{(i-2)!j!} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_0^1 \int_0^1 \right. \\
&\quad \left(\frac{y_2}{y_3} \right)^{-(\alpha(1-t_2)+\beta t_2)t_1} y_2^{\gamma x_1 - (\alpha(1-t_2)+\beta t_2)t_1 x_1} y_3^{\alpha x_2 + \beta x_3} (1-t_1)^{i-2} t_1 \\
&\quad \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(\frac{y_3}{n} \right)^{-(\alpha(1-t_2)+\beta t_2)t_1} \left(x_1 + \frac{\log y_2/n}{\log y_2} \right)^i \left(x_2 + x_3 + \frac{\log y_3/n}{\log y_3} \right)^j dt_1 dt_2 \Big]_{\underline{x}=0} \\
&\quad + O_\varepsilon \left(\mathcal{L}^{i-2+\varepsilon} \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(\frac{y_3}{n} \right)^{-\nu} \right) + O(\mathcal{L}^{i+j-3}).
\end{aligned}$$

Using Lemmas 4.1 and 4.2, the O -terms are $\ll_\varepsilon \mathcal{L}^{i+1+\varepsilon} + \mathcal{L}^{i+j-3}$, while the first term is

$$\begin{aligned}
&\frac{2(\log y_2)^{i-2} (\log y_3)^j}{3(i-2)!j!} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_0^1 \int_0^1 \int_0^1 \right. \\
&\quad y_2^{\gamma x_1 - (\alpha(1-t_2)+\beta t_2)t_1(1+x_1)} y_3^{\alpha x_2 + \beta x_3 + (\alpha(1-t_2)+\beta t_2)t_1(1-u)} (1-t_1)^{i-2} t_1 (1-u)^3
\end{aligned}$$

$$\left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right)^i (x_2 + x_3 + u)^j dt_1 dt_2 du \Big|_{\underline{x}=0} + O(\mathcal{L}^{i+j-3}).$$

Putting this back to (28) and using the assumption that $j \geq 4$ we get

$$\begin{aligned} I_{23}^1 &= \frac{2\widehat{w}(0)}{3(\log y_2)^2} \zeta(1+\alpha+\gamma) \zeta(1+\beta+\gamma) \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_0^1 \int_0^1 \int_0^1 \right. \\ &\quad \frac{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 (1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1 (1-u)}{y_3} t_1 \left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right)^2 (1-u)^3 \\ &\quad \left. P_2'' \left(\left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right) (1-t_1) \right) P_3(x_2 + x_3 + u) dt_1 dt_2 du \right]_{\underline{x}=0} + O_\varepsilon(\mathcal{L}^{-1+\varepsilon}). \end{aligned}$$

6.2. Deduction of Lemma 3.2. Combining I_{23}^1 , I_{23}^2 and I_{23}^3 we have

$$\begin{aligned} I_{23} &= \frac{2\widehat{w}(0)}{3(\log y_2)^2} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_0^1 \int_0^1 \int_0^1 U_2(\underline{x}) t_1 \left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right)^2 (1-u)^3 \right. \\ &\quad \left. P_2'' \left(\left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2}(1-u) \right) (1-t_1) \right) P_3(x_2 + x_3 + u) dt_1 dt_2 du \right]_{\underline{x}=0} + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}), \end{aligned}$$

where

$$\begin{aligned} U_2 &= \frac{\frac{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 (1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1 (1-u)}{y_3}}{(\alpha + \gamma)(\beta + \gamma)} \\ &\quad - \frac{T^{-(\alpha+\gamma)} y_2^{-\alpha x_1 - (-\gamma(1-t_2) + \beta t_2) t_1 (1+x_1)} y_3^{-\gamma x_2 + \beta x_3 + (-\gamma(1-t_2) + \beta t_2) t_1 (1-u)}}{(\alpha + \gamma)(\beta - \alpha)} \\ &\quad - \frac{T^{-(\beta+\gamma)} y_2^{-\beta x_1 - (\alpha(1-t_2) - \gamma t_2) t_1 (1+x_1)} y_3^{\alpha x_2 - \gamma x_3 + (\alpha(1-t_2) - \gamma t_2) t_1 (1-u)}}{(\alpha - \beta)(\beta + \gamma)}. \end{aligned}$$

As in the previous section we can write

$$\begin{aligned} U_2 &= \frac{\frac{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 (1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1 (1-u)}{y_3}}{\beta - \alpha} \\ &\quad \left(\frac{1 - (T y_2^{x_1 - t_1(1-t_2)(1+x_1)} y_3^{x_2 + t_1(1-t_2)(1-u)})^{-(\alpha+\gamma)}}{\alpha + \gamma} \right) \\ &\quad - \frac{\frac{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 (1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1 (1-u)}{y_3}}{\beta - \alpha} \\ &\quad \left(\frac{1 - (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_3 + t_1 t_2(1-u)})^{-(\beta+\gamma)}}{\beta + \gamma} \right). \end{aligned}$$

Applying (25) leads to

$$\begin{aligned} U_2 &= \frac{\mathcal{L}}{\beta - \alpha} \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1(1-t_2)(\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) \\ &\quad \frac{\gamma x_1 - (\alpha(1-t_2) + \beta t_2) t_1 (1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1 (1-u)}{y_3} \\ &\quad \int_0^1 (T y_2^{x_1 - t_1(1-t_2)(1+x_1)} y_3^{x_2 + t_1(1-t_2)(1-u)})^{-(\alpha+\gamma) t_3} dt_3 \\ &\quad - \frac{\mathcal{L}}{\beta - \alpha} \left(1 + \vartheta_2 x_1 + \vartheta_3 x_3 - t_1 t_2 (\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) \end{aligned}$$

$$\frac{\gamma_{x_1 - (\alpha(1-t_2) + \beta t_2)} t_1(1+x_1)}{y_2} \frac{\alpha x_2 + \beta x_3 + (\alpha(1-t_2) + \beta t_2) t_1(1-u)}{y_3} \int_0^1 (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_3 + t_1 t_2(1-u)})^{-(\beta+\gamma)t_3} dt_3.$$

Changing $t_2 \rightarrow 1 - t_2$ in the first term, and changing the roles of the variables x_2 with x_3 in the second term yields

$$\begin{aligned} I_{23} = & \frac{2\widehat{w}(0)\mathcal{L}}{3(\log y_2)^2} \frac{d^6}{dx_1^2 dx_2^2 dx_3^2} \left[\int_{[0,1]^4} V_2(\underline{x}) \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2 (\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) \right. \\ & t_1 \left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2} (1-u) \right)^2 (1-u)^3 P_2'' \left(\left(x_1 + 1 - \frac{\vartheta_3}{\vartheta_2} (1-u) \right) (1-t_1) \right) \\ & \left. P_3(x_2 + x_3 + u) dt_1 dt_2 dt_3 du \right]_{\underline{x}=0} + O_\varepsilon(T\mathcal{L}^{-1+\varepsilon}), \end{aligned}$$

where

$$\begin{aligned} V_2 = & \frac{1}{\beta - \alpha} \left(y_2^{\gamma_{x_1 - (\alpha t_2 + \beta(1-t_2))} t_1(1+x_1)} y_3^{\alpha x_2 + \beta x_3 + (\alpha t_2 + \beta(1-t_2)) t_1(1-u)} \right. \\ & (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{-(\alpha+\gamma)t_3} \\ & - y_2^{\gamma_{x_1 - (\alpha(1-t_2) + \beta t_2)} t_1(1+x_1)} y_3^{\alpha x_3 + \beta x_2 + (\alpha(1-t_2) + \beta t_2) t_1(1-u)} \\ & \left. (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{-(\beta+\gamma)t_3} \right). \end{aligned}$$

Using (25) again implies that V_2 equals

$$\begin{aligned} & y_2^{\gamma_{x_1 - (\alpha t_2 + \beta(1-t_2))} t_1(1+x_1)} y_3^{\alpha x_2 + \beta x_3 + (\alpha t_2 + \beta(1-t_2)) t_1(1-u)} (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{-(\alpha+\gamma)t_3} \\ & \left(\frac{1 - \left(y_2^{t_1(2t_2-1)(1+x_1)} y_3^{-x_2+x_3-t_1(2t_2-1)(1-u)} (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{t_3} \right)^{-(\beta-\alpha)}}{\beta - \alpha} \right) \\ = & \mathcal{L} \left(-\vartheta_3(x_2 - x_3) + t_1(2t_2 - 1)(\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right. \\ & \left. + t_3 \left(1 + \vartheta_2 x_1 + \vartheta_3 x_2 - t_1 t_2 (\vartheta_2(1+x_1) - \vartheta_3(1-u)) \right) \right) \\ & y_2^{\gamma_{x_1 - (\alpha t_2 + \beta(1-t_2))} t_1(1+x_1)} y_3^{\alpha x_2 + \beta x_3 + (\alpha t_2 + \beta(1-t_2)) t_1(1-u)} (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{-(\alpha+\gamma)t_3} \\ & \int_0^1 \left(y_2^{t_1(2t_2-1)(1+x_1)} y_3^{-x_2+x_3-t_1(2t_2-1)(1-u)} (T y_2^{x_1 - t_1 t_2(1+x_1)} y_3^{x_2 + t_1 t_2(1-u)})^{t_3} \right)^{-(\beta-\alpha)t_4} dt_4, \end{aligned}$$

and the lemma follows.

7. PROOF OF LEMMA 3.1

7.1. Reduction to a contour integral. We first state the twisted fourth moment of the Riemann zeta-function [2].

Theorem 7.1 (Bettin, Bui, Li and Radziwiłł). *Suppose $H, K \leq T^{1/4-\varepsilon}$. Then we have*

$$\sum_{\substack{h \leq H \\ k \leq K}} \frac{a_h \overline{a_k}}{\sqrt{hk}} \int_{-\infty}^{\infty} \zeta\left(\frac{1}{2} + \alpha + it\right) \zeta\left(\frac{1}{2} + \beta + it\right) \zeta\left(\frac{1}{2} + \gamma - it\right) \zeta\left(\frac{1}{2} + \delta - it\right) \left(\frac{h}{k}\right)^{it} w(t) dt$$

$$\begin{aligned}
&= \sum_{\substack{h \leq H \\ k \leq K}} \frac{a_h \overline{a_k}}{\sqrt{hk}} \int_{-\infty}^{\infty} w(t) \left\{ Z_{\alpha, \beta, \gamma, \delta}(h, k) + \left(\frac{t}{2\pi} \right)^{-(\alpha+\gamma)} Z_{-\gamma, \beta, -\alpha, \delta}(h, k) \right. \\
&\quad + \left(\frac{t}{2\pi} \right)^{-(\alpha+\delta)} Z_{-\delta, \beta, \gamma, -\alpha}(h, k) + \left(\frac{t}{2\pi} \right)^{-(\beta+\gamma)} Z_{\alpha, -\gamma, -\beta, \delta}(h, k) \\
&\quad \left. + \left(\frac{t}{2\pi} \right)^{-(\beta+\delta)} Z_{\alpha, -\delta, \gamma, -\beta}(h, k) + \left(\frac{t}{2\pi} \right)^{-(\alpha+\beta+\gamma+\delta)} Z_{-\gamma, -\delta, -\alpha, -\beta}(h, k) \right\} dt + O_\varepsilon(T^{1-\varepsilon})
\end{aligned}$$

uniformly for $\alpha, \beta, \gamma, \delta \ll \mathcal{L}^{-1}$, where

$$Z_{\alpha, \beta, \gamma, \delta}(h, k) = \sum_{kab=hcd} \frac{1}{a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma} d^{1/2+\delta}}.$$

Recall that $I_3(\alpha, \beta, \gamma, \delta)$ is defined by (10). We write

$$I_3(\alpha, \beta, \gamma, \delta) = I_3^1 + I_3^2 + I_3^3 + I_3^4 + I_3^5 + I_3^6 + O_\varepsilon(T^{1-\varepsilon})$$

correspondingly to the decomposition in Theorem 7.1. We first work on I_3^1 , which is equal to

$$\widehat{w}(0) \sum_{l, m} \frac{\mu_2(m) \mu_2(l) P_3\left(\frac{\log y_3/m}{\log y_3}\right) P_3\left(\frac{\log y_3/l}{\log y_3}\right)}{\sqrt{lm}} \sum_{mab=lcd} \frac{1}{a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma} d^{1/2+\delta}}.$$

In view of (19) we get

$$\begin{aligned}
I_3^1 &= \widehat{w}(0) \sum_{i, j} \frac{c_i c_j i! j!}{(\log y_3)^{i+j}} \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} y_3^{u+v} \\
&\quad \sum_{mab=lcd} \frac{\mu_2(m) \mu_2(l)}{m^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma} d^{1/2+\delta}} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}.
\end{aligned}$$

The arithmetical sum is

$$\begin{aligned}
&\sum_{mab=lcd} \frac{\mu_2(m) \mu_2(l)}{m^{1/2+u} l^{1/2+v} a^{1/2+\alpha} b^{1/2+\beta} c^{1/2+\gamma} d^{1/2+\delta}} \\
&= C(\alpha, \beta, \gamma, \delta, u, v) \frac{\zeta(1+\alpha+\gamma) \zeta(1+\alpha+\delta) \zeta(1+\beta+\gamma) \zeta(1+\beta+\delta) \zeta(1+u+v)^4}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+\delta+u)^2},
\end{aligned} \tag{33}$$

where $C(\alpha, \beta, \gamma, \delta, u, v)$ is an arithmetical factor converging absolutely in a product of half-planes containing the origin. Hence

$$I_3^1 = \widehat{w}(0) \zeta(1+\alpha+\gamma) \zeta(1+\alpha+\delta) \zeta(1+\beta+\gamma) \zeta(1+\beta+\delta) \sum_{i, j} \frac{c_i c_j i! j!}{(\log y_3)^{i+j}} L_{i, j}, \tag{34}$$

where

$$L_{i, j} = \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} y_3^{u+v} \frac{C(\alpha, \beta, \gamma, \delta, u, v) \zeta(1+u+v)^4}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+\delta+u)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}.$$

Using the Dirichlet series for $\zeta(1+u+v)^4$ and reversing the order of summation and integration, we have

$$\begin{aligned}
L_{i, j} &= \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(\frac{1}{2\pi i} \right)^2 \int_{(1)} \int_{(1)} \left(\frac{y_3}{n} \right)^{u+v} \\
&\quad \frac{C(\alpha, \beta, \gamma, \delta, u, v)}{\zeta(1+\alpha+v)^2 \zeta(1+\beta+v)^2 \zeta(1+\gamma+u)^2 \zeta(1+\delta+u)^2} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}. \tag{35}
\end{aligned}$$

Here we are able to restrict the sum over n to $n \leq y_3$ by moving the u, v -integrals far to the right. We now move the contours of integration to $\text{Re}(u) = \text{Re}(v) \asymp \mathcal{L}^{-1}$. Bounding the

integrals trivially shows that $L_{i,j} \ll \mathcal{L}^{i+j-4}$. As before we can replace $C(\alpha, \beta, \gamma, \delta, u, v)$ by $C(0, 0, 0, 0, 0, 0)$ in $L_{i,j}$ with an error of size $O(\mathcal{L}^{i+j-5})$. By letting $\alpha = \beta = \gamma = \delta = u = v = s$ in (33), it is easy to verify that $C(0, 0, 0, 0, 0, 0) = 1$. The u and v variables in (35) are now separated so that

$$L_{i,j} = \sum_{n \leq y_3} \frac{d_4(n)}{n} N_i(\gamma, \delta) N_j(\alpha, \beta) + O(\mathcal{L}^{i+j-5}),$$

where the function $N_j(\alpha, \beta)$ is defined in (31). Using Lemma 6.2 we obtain

$$\begin{aligned} L_{i,j} &= \frac{(\log y_3)^{i+j-8}}{i!j!} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} \right. \\ &\quad \left. \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(x_3 + x_4 + \frac{\log y_3/n}{\log y_3} \right)^i \left(x_1 + x_2 + \frac{\log y_3/n}{\log y_3} \right)^j \right]_{\underline{x}=0} \\ &\quad + O_\varepsilon \left((\mathcal{L}^{i-4+\varepsilon} + \mathcal{L}^{j-4+\varepsilon}) \sum_{n \leq y_3} \frac{d_4(n)}{n} \left(\frac{y_3}{n} \right)^{-\nu} \right) + O(\mathcal{L}^{i+j-5}). \end{aligned}$$

In view of Lemmas 4.1 and 4.2, the O -terms are $\ll_\varepsilon \mathcal{L}^{i-1+\varepsilon} + \mathcal{L}^{j-1+\varepsilon} + \mathcal{L}^{i+j-5}$, while the first term is

$$\begin{aligned} &\frac{(\log y_3)^{i+j-4}}{6i!j!} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[\int_0^1 y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (1-u)^3 (x_3 + x_4 + u)^i (x_1 + x_2 + u)^j du \right]_{\underline{x}=0} \\ &\quad + O(\mathcal{L}^{i+j-5}). \end{aligned}$$

Combining this with (34), and using the assumption $i, j \geq 4$, we get

$$\begin{aligned} I_3^1 &= \frac{\widehat{w}(0)}{6(\log y_3)^4} \zeta(1+\alpha+\gamma) \zeta(1+\alpha+\delta) \zeta(1+\beta+\gamma) \zeta(1+\beta+\delta) \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[\int_0^1 \right. \\ &\quad \left. y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (1-u)^3 P_3(x_1 + x_2 + u) P_3(x_3 + x_4 + u) du \right]_{\underline{x}=0} + O_\varepsilon(\mathcal{L}^{-1+\varepsilon}). \end{aligned}$$

7.2. Deduction of Lemma 3.1. Recall that I_3^2 is essentially obtained by multiplying I_3^1 with $T^{-(\alpha+\gamma)}$ and changing the shifts $\alpha \leftrightarrow -\gamma$, I_3^3 is obtained by multiplying I_3^1 with $T^{-(\alpha+\delta)}$ and changing the shifts $\alpha \leftrightarrow -\delta$, I_3^4 is obtained by multiplying I_3^1 with $T^{-(\beta+\gamma)}$ and changing the shifts $\beta \leftrightarrow -\gamma$, I_3^5 is obtained by multiplying I_3^1 with $T^{-(\beta+\delta)}$ and changing the shifts $\beta \leftrightarrow -\delta$, and I_3^6 is obtained by multiplying I_3^1 with $T^{-(\alpha+\beta+\gamma+\delta)}$ and changing the shifts $\alpha \leftrightarrow -\gamma$ and $\beta \leftrightarrow -\delta$. Hence

$$\begin{aligned} I_3(\alpha, \beta, \gamma, \delta) &= \frac{\widehat{w}(0)}{6(\log y_3)^4} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[\int_0^1 \right. \\ &\quad \left. U(\underline{x})(1-u)^3 P_3(x_1 + x_2 + u) P_3(x_3 + x_4 + u) du \right]_{\underline{x}=0} + O_\varepsilon(\mathcal{L}^{-1+\varepsilon}), \end{aligned} \tag{36}$$

where

$$\begin{aligned} U &= \frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(\alpha + \gamma)(\alpha + \delta)(\beta + \gamma)(\beta + \delta)} - \frac{T^{-(\alpha+\gamma)} y_3^{-\gamma x_1 + \beta x_2 - \alpha x_3 + \delta x_4}}{(\alpha + \gamma)(-\gamma + \delta)(\beta - \alpha)(\beta + \delta)} \\ &\quad - \frac{T^{-(\alpha+\delta)} y_3^{-\delta x_1 + \beta x_2 + \gamma x_3 - \alpha x_4}}{(-\delta + \gamma)(\alpha + \delta)(\beta + \gamma)(\beta - \alpha)} - \frac{T^{-(\beta+\gamma)} y_3^{\alpha x_1 - \gamma x_2 - \beta x_3 + \delta x_4}}{(\alpha - \beta)(\alpha + \delta)(\beta + \gamma)(-\gamma + \delta)} \\ &\quad - \frac{T^{-(\beta+\delta)} y_3^{\alpha x_1 - \delta x_2 + \gamma x_3 - \beta x_4}}{(\alpha + \gamma)(\alpha - \beta)(-\delta + \gamma)(\beta + \delta)} + \frac{T^{-(\alpha+\beta+\gamma+\delta)} y_3^{-\gamma x_1 - \delta x_2 - \alpha x_3 - \beta x_4}}{(\alpha + \gamma)(\gamma + \beta)(\delta + \alpha)(\beta + \delta)}. \end{aligned}$$

We write

$$\frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(\alpha + \gamma)(\alpha + \delta)(\beta + \gamma)(\beta + \delta)} = \frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(\alpha + \gamma)(-\gamma + \delta)(\beta - \alpha)(\beta + \delta)} - \frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(-\gamma + \delta)(\alpha + \delta)(\beta + \gamma)(\beta - \alpha)}$$

and

$$\frac{T^{-(\alpha + \beta + \gamma + \delta)} y_3^{-\gamma x_1 - \delta x_2 - \alpha x_3 - \beta x_4}}{(\alpha + \gamma)(\alpha + \delta)(\beta + \gamma)(\beta + \delta)} = \frac{T^{-(\alpha + \beta + \gamma + \delta)} y_3^{-\gamma x_1 - \delta x_2 - \alpha x_3 - \beta x_4}}{(\alpha + \gamma)(-\gamma + \delta)(\beta - \alpha)(\beta + \delta)} - \frac{T^{-(\alpha + \beta + \gamma + \delta)} y_3^{-\gamma x_1 - \delta x_2 - \alpha x_3 - \beta x_4}}{(-\gamma + \delta)(\alpha + \delta)(\beta + \gamma)(\beta - \alpha)}. \quad (37)$$

Notice that we can change the roles of x_1 with x_2 , or of x_3 with x_4 in any term of $U(\underline{x})$ without affecting the value of $I_3(\alpha, \beta, \gamma, \delta)$ in (36). Applying both changes to the last term in (37), we can replace $U(\underline{x})$ with

$$\begin{aligned} & \frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(-\gamma + \delta)(\beta - \alpha)} \left(\frac{1 - (T y_3^{x_1 + x_3})^{-(\alpha + \gamma)}}{\alpha + \gamma} \right) \left(\frac{1 - (T y_3^{x_2 + x_4})^{-(\beta + \delta)}}{\beta + \delta} \right) \\ & - \frac{y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4}}{(-\gamma + \delta)(\beta - \alpha)} \left(\frac{1 - (T y_3^{x_1 + x_4})^{-(\alpha + \delta)}}{\alpha + \delta} \right) \left(\frac{1 - (T y_3^{x_2 + x_3})^{-(\beta + \gamma)}}{\beta + \gamma} \right). \end{aligned}$$

Using (25) we then get

$$\begin{aligned} I_3(\alpha, \beta, \gamma, \delta) &= \frac{\mathcal{L}^2 \hat{w}(0)}{6(\log y_3)^4} \frac{d^8}{dx_1^2 dx_2^2 dx_3^2 dx_4^2} \left[\int_{[0,1]^3} \frac{V_1(\underline{x}, t_1, t_2) - V_2(\underline{x}, t_1, t_2)}{(-\gamma + \delta)(\beta - \alpha)} \right. \\ & \quad \left. (1 - u)^3 P_3(x_1 + x_2 + u) P_3(x_3 + x_4 + u) dt_1 dt_2 du \right]_{\underline{x}=0} + O_\varepsilon(\mathcal{L}^{-1+\varepsilon}), \end{aligned}$$

where

$$V_1 = y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (T y_3^{x_1 + x_3})^{-(\alpha + \gamma) t_1} (T y_3^{x_2 + x_4})^{-(\beta + \delta) t_2} (1 + \vartheta_3(x_1 + x_3)) (1 + \vartheta_3(x_2 + x_4))$$

and

$$V_2 = y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (T y_3^{x_1 + x_4})^{-(\alpha + \delta) t_1} (T y_3^{x_2 + x_3})^{-(\beta + \gamma) t_2} (1 + \vartheta_3(x_1 + x_4)) (1 + \vartheta_3(x_2 + x_3)).$$

Again notice that $I_3(\alpha, \beta, \gamma, \delta)$ is unchanged if we swap any of these pairs of variables $x_1 \leftrightarrow x_2$, $x_3 \leftrightarrow x_4$ and $t_1 \leftrightarrow t_2$ in $V_1(\underline{x}, t_1, t_2)$ or $V_2(\underline{x}, t_1, t_2)$. We next replace $V_1 - V_2$ in the integrand with

$$\begin{aligned} & \frac{1}{2} \left(V_1(x_1, x_2, x_3, x_4, t_1, t_2) - V_2(x_2, x_1, x_3, x_4, t_2, t_1) - V_2(x_1, x_2, x_4, x_3, t_1, t_2) \right. \\ & \quad \left. + V_1(x_2, x_1, x_4, x_3, t_2, t_1) \right), \end{aligned}$$

which is

$$\begin{aligned} & \frac{(1 + \vartheta_3(x_1 + x_3))(1 + \vartheta_3(x_2 + x_4))}{2} \left(y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} (T y_3^{x_1 + x_3})^{-(\alpha + \gamma) t_1} (T y_3^{x_2 + x_4})^{-(\beta + \delta) t_2} \right. \\ & \quad - y_3^{\alpha x_2 + \beta x_1 + \gamma x_3 + \delta x_4} (T y_3^{x_2 + x_4})^{-(\alpha + \delta) t_2} (T y_3^{x_1 + x_3})^{-(\beta + \gamma) t_1} \\ & \quad - y_3^{\alpha x_1 + \beta x_2 + \gamma x_4 + \delta x_3} (T y_3^{x_1 + x_3})^{-(\alpha + \delta) t_1} (T y_3^{x_2 + x_4})^{-(\beta + \gamma) t_2} \\ & \quad \left. + y_3^{\alpha x_2 + \beta x_1 + \gamma x_4 + \delta x_3} (T y_3^{x_2 + x_4})^{-(\alpha + \gamma) t_2} (T y_3^{x_1 + x_3})^{-(\beta + \delta) t_1} \right) \\ &= \frac{(1 + \vartheta_3(x_1 + x_3))(1 + \vartheta_3(x_2 + x_4))}{2} y_3^{\alpha x_1 + \beta x_2 + \gamma x_3 + \delta x_4} \end{aligned}$$

$$(Ty_3^{x_1+x_3})^{-(\alpha+\gamma)t_1}(Ty_3^{x_2+x_4})^{-(\beta+\delta)t_2}\left(1-\left(T^{t_1-t_2}y_3^{-x_1+x_2+(x_1+x_3)t_1-(x_2+x_4)t_2}\right)^{-(\beta-\alpha)}\right) \\ \left(1-\left(T^{t_1-t_2}y_3^{-x_3+x_4+(x_1+x_3)t_1-(x_2+x_4)t_2}\right)^{-(\delta-\gamma)}\right).$$

Using (25) again and simplifying we obtain Lemma 3.1.

REFERENCES

- [1] R. Balasubramanian, J. B. Conrey, D. R. Heath-Brown, *Asymptotic mean square of the product of the Riemann zeta-function and a Dirichlet polynomial*, J. Reine Angew. Math. **357** (1985), 161–181.
- [2] S. Bettin, H. M. Bui, X. Li, M. Radziwiłł, preprint.
- [3] H. M. Bui, J. B. Conrey, M. P. Young, *More than 41% of the zeros of the zeta function are on the critical line*, Acta Arith. **150** (2011), 35–64.
- [4] J. B. Conrey, *More than two fifths of the zeros of the Riemann zeta function are on the critical line*, J. Reine Angew. Math. **399** (1989), 1–26.
- [5] J. B. Conrey, H. Iwaniec, K. Soundararajan, *Critical zeros of Dirichlet L-functions*, J. Reine Angew. Math. **681** (2013), 175–198.
- [6] J. B. Conrey, N. C. Snaith, *Applications of the L-functions ratios conjectures*, Proc. Lond. Math. Soc. **94** (2007), 594–646.
- [7] S. Feng, *Zeros of the Riemann zeta function on the critical line*, J. Number Theory **132** (2012), 511–542.
- [8] S. Feng, *Zeros of the Riemann zeta function on the critical line*, arXiv <http://arxiv.org/abs/1003.0059>
- [9] D. R. Heath-Brown, *Simple zeros of the Riemann zeta function on the critical line*, Bull. London Math. Soc. **11** (1979), 17–18.
- [10] N. Levinson, *More than one third of the zeros of Riemann’s zeta function are on $\sigma = 1/2$* . Adv. Math. **13** (1974), 383–436.
- [11] A. Selberg, *On the zeros of Riemann’s zeta-function*, Skr. Norske Vid. Akad. Oslo I. (1942), no. 10, 59 pp.
- [12] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, revised by D. R. Heath-Brown, Clarendon Press, second edition, 1986.

SCHOOL OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL BS8 1TW, UK
E-mail address: hung.bui@bristol.ac.uk